

Subject Index

AACVPR. *See* American Association of Cardiovascular and Pulmonary Rehabilitation
Acute coronary syndrome, high-risk non-ST elevation, 236
Acute exacerbations of COPD, community pulmonary rehabilitation after hospitalization for, 186
Adherence
  lived experience of, in heart failure patients, 158
  qualitative research on, 164
Aerobic capacity, peak, entering cardiac rehabilitation, 293
Aerobic-resistance training
  effects on diminution of skeletal muscle fatigue in men with coronary heart disease, 312
  women's self-efficacy after, in cardiac rehabilitation, 310
Aerobic training
  and β-blockers, 354
  women's self-efficacy after, in cardiac rehabilitation, 310
Age
  advanced, comprehensive cardiac rehabilitation program and cardiac-related events in, 302
  and quality of life after cardiac rehabilitation, 304
Airway obstruction, severe chronic, 236
American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR)
  20th anniversary, 1, 241
  2005 Annual Meeting, 288
  2005 Annual Meeting abstracts, 290
  Award of Excellence lecture, 67
  first 20 years, 242
  position statement on depression screening in cardiac rehabilitation, 11
Atkins diet for weight loss and heart disease risk reduction, 184
Auditory stimuli, distractive, 50

Behavioral management
  enhancing patient motivation for change, 207
  of heart failure, 164
β-blockers, exercise training and, 354
Blood
  capillary vs venous insulin responses, 305
  glucose levels during exercise stress testing, 222
BMI. *See* Body mass index
Body composition, whole-body exercise training effects on, 124
Body mass index, exercise capacity gains and, 303
Borg's Rating of Perceived Exertion, 309
Bruce treadmill protocol
  comparison with modified Bruce protocol, 290
  modified, 290
Bypass surgery, coronary
  cognitive performance and psychosocial and physical functioning after, 300
  controlled breathing program and heart rate variability after, 308
  off-pump versus on-pump cardiac rehabilitation outcomes for, 294
  ST-segment depression after, during phase II cardiac rehabilitation, 297

C-reactive protein
  cardiac rehabilitation and, in cardiovascular disease, 332
  cholesterol and, 304
  energetic composition of prior meal and acute exercise and, in type 2 diabetics, 310
Canadian Association of Cardiac Rehabilitation: 15th Annual Meeting and Symposium abstracts, 310
Capillary blood insulin response, 305
Carbon dioxide, resting end-tidal ($P_{ET}CO2$), 177
Cardiac rehabilitation
  AACVPR position statement on depression screening in, 11
  adherence to
    exercise capacity and health-related quality of life and, 291
    social support, family functioning, depression, and quality of life and, 300
  age and quality of life after, 304
  C-reactive protein and, in cardiovascular disease, 332
  case for, 249
  cognitive performance and, in older adults, 173
  complications during, in France, 296
  depressive disorders in men and women enrolled in, 71
  in diabetes, 59
  effort capacity during third month of, 210
  enrollment factors, 343
  exercise capacity gains and BMI in, 303
  exercise capacity 2 years after, 14
  first 20 years, 243
  geographic distribution of programs, 80
  home-based
    patient preferences for, 24
    as preferred and viable option, 30
  horticultural therapy effects on mood and heart rate in, 270
  hospital-based: patient preferences for, 24
  lipid outcomes one year after, 295
  lipids and, 257, 264
  long-term effects of, 103
  in metabolic syndrome, 59
  in metabolic syndrome and diabetes mellitus, 298
  motivational interviewing and secondary prevention gains in, 200
  nonparticipation in, 88
  in obesity, 62
  occupational outcomes in, 311
  in off-pump versus on-pump coronary bypass surgery, 294
  paradigms of, 103
  patient outcome indicators, Delphi study, 303
  peak aerobic capacity entering, 293
  phase II
    effects on depression and quality of life, 296
    gender differences in risk factor status and completion of, 304
    pre and post predicted VO2 max measurements in, 303
    screening for depression in, 299
    ST-segment depression post-CABG during, 297
  physical activity 2 years after, 14

quality of life outcomes
and age, 304
in elderly, 295
risk factor status on entry into, male versus female, 297
secondary prevention gains in, 200
strength assessment with hand-held dynamometry in, 266
Telehealth Program pilot (Texas State University), 302
Transtheoretical Model-based staff training and clinical intervention in, 301
2 year follow-up after, 14
utilization of services, 350
women's self-efficacy after aerobic or aerobic-resistance training in, 310
Cardiac rehabilitation programs
comprehensive
effects on cardiac-related events in advanced age, 302
effects on cardiac-related events in female cohort, 294
distribution of, 85
outpatient: medical director responsibilities for, 315
Cardiac-related events: comprehensive cardiac rehabilitation program and, in female cohort, 294
Cardiac risk: depression and, 78
Cardiological treatments: life-years gained from, in England and Wales, 185
Cardiopulmonary exercise testing in heart failure, 292
Cardiopulmonary rehabilitation: and depression assessment, 291
Cardiovascular care: quality indicators in, 249
Cardiovascular disease: cardiac rehabilitation and C-reactive protein in, 332
Cardiovascular events, exercise-related, 189
Cardiovascular function: sitting posture and, in COPD, 305
Cardiovascular Health Best Practice Project, 311
Cardiovascular maintenance: lipid outcomes in, one year after cardiac rehabilitation, 295
Cardiovascular risk markers: exercise and stress management training effects on, in ischemic heart disease, 235
Caregivers, spouse: spirituality and religiosity and perception of quality of life and confidence, 299
Cholesterol: C-reactive protein and, 304
Chronic airway obstruction, severe: ventilatory support during training in, 236
Chronic heart failure
relaxation response randomized trial of, 149
resting end-tidal carbon dioxide in, reliability of, 177
Chronic obstructive pulmonary disease
combination tiotropium and pulmonary rehabilitation in, 386
community management of, 378
community pulmonary rehabilitation after hospitalization for acute exacerbations of, 186
daily activity profile in, 181
distractive auditory stimuli on upper extremity training in, 50
endurance and strength training in patients with, 123
exercise outcomes after pulmonary rehabilitation in, 186
exercise training and dyspnea self-management program in, 275
home-based pulmonary rehabilitation in, 378
Human Activity Profile usefulness in, 115
incremental treadmill walking and skeletal muscle oxygen saturation in, 298
metabolic syndrome in, 226, 233
muscle function and shuttle walking performance in, 43
PUFAs and exercise capacity in, 237
quality of life outcome measures in, 321
Seattle Obstructive Lung Disease Questionnaire (SOLDQ), 107
sitting posture and pulmonary and cardiovascular function in, 305
strength training in patients with, 123
whole-body exercise training effects on body composition and functional capacity in, 124
Cognitive function: physical activity and, in older women, 56
Cognitive performance
cardiac rehabilitation and, in older adults, 173
and psychosocial and physical functioning after coronary artery bypass, 300
Community management of COPD, 378
Community pulmonary rehabilitation: after hospitalization for acute exacerbations of COPD, 186
Comorbidity indexes: and classification of CAD, 293
Compliance
lived experience of adherence for heart failure patients, 158
qualitative research on, 164
Comprehensive cardiac rehabilitation programs
effects on cardiac-related events, in female cohort, 294
effects on cardiac-related events in advanced age, 302
Computed tomography, coronary, 3
Control, perceived: and PTSD after myocardial infarction, 166
COPD. *See* Chronic obstructive pulmonary disease
Coronary artery bypass graft surgery
cognitive performance and psychosocial and physical functioning after, 300
controlled breathing program and heart rate variability after, 308
off-pump versus on-pump cardiac rehabilitation outcomes for, 294
pain or difficulty ratings as indicators of functional deficits after, 300
ST-segment depression after, during phase II cardiac rehabilitation, 297
Coronary artery calcification: stress-induced myocardial ischemia and, 305
Coronary artery disease
cardiac rehabilitation and lipids in, 257
comorbidity indexes and classification of, 293
maximal graded exercise testing and glutathione in, 292
Coronary computed tomography, in coronary risk assessment, 3
Coronary heart disease
aerobic-resistance training and skeletal muscle fatigue in men with, 312
causal attribution for, in female patients, 135
women's perceptions of, 146
Coronary risk assessment, 3
Coronary risk factors, 305
Counting Talk Test
comparison with heart rate measures and Bork's Rating of Perceived Exertion, 309
comparison with oxygen consumption, 308
CT. *See* Computed tomography
Cycle exercise. *See* North Carolina 6-minute cycle test

Daily activity profile in COPD, 181
Depression
in cardiac rehabilitation enrollees, 71
and cardiac risk, 78
and heart rate recovery after treadmill stress testing, 307
and medical adherence in cardiac rehabilitation, 300
and phase II cardiac rehabilitation, 296
screening for, in phase II cardiac rehabilitation, 299
Depression screening
in cardiac rehabilitation: AACVPR position statement on, 11
in cardiopulmonary rehabilitation, 291
Diabetes mellitus
cardiac rehabilitation and exercise training in, 59
cardiac rehabilitation and metabolic syndrome in, 298
fitness in, 59
high-risk asymptomatic candidates for screening stress SPECT imaging, 122
type 2: energetic composition of prior meal and acute exercise and C-reactive protein in, 310
Diastolic heart failure, pedometer steps in, 293
Diets for weight loss and heart disease risk reduction, 184
Difficulty ratings: and functional deficits after coronary artery bypass, 300
Disabilities, persons with: obstacles to exercise and rehabilitation equipment use for, 301
Distractive auditory stimuli on upper extremity training, in COPD, 50

Dynamometry, hand-held, 266
Dyspnea
distractive auditory stimuli during upper extremity training and, in COPD, 50
exercise training and self-management of, 275, 286

Effort capacity during third month of cardiac rehabilitation, 210
8 week outpatient pulmonary rehabilitation program, 296
Elderly
cardiac rehabilitation patients: quality of life outcomes for, 295
obese, with osteoarthritis of knee: home-based walking and strength training program for, 309
Electrocardiography, exercise: diagnostic value of heart rate during, 127
End-tidal carbon dioxide, resting ($P_{ET}CO2$): reliability in chronic heart failure, 177
Endurance training: in COPD, 123
Energetic composition: and C-reactive protein in type 2 diabetics, 310
England and Wales: life-years gained from cardiological treatments and risk factor changes in, 185
Exercise
acute
and C-reactive protein in type 2 diabetics, 310
and quality of life, in heart failure, 311
anti-inflammatory effects of, 339
capacity for
adherence to cardiac rehabilitation and, 291
and body mass index, in cardiac rehabilitation, 303
and depression and heart rate recovery after treadmill stress testing, 307
effort capacity during third month of cardiac rehabilitation, 210
nomogram prognostic value for, in women, 386
and polyunsaturated fatty acids, in COPD, 237
and quadriceps weakness, in idiopathic pulmonary fibrosis, 386
2 years after cardiac rehabilitation, 14
and whole-body exercise training, in COPD, 124
after cardiac hospitalization: Tracking Exercise After Cardiac Hospitalization (TEACH) Study, 311
cardiovascular events associated with, 189
and cardiovascular risk markers, in ischemic heart disease, 235
duration of: and postexercise hypotension, 366
and enrollment in cardiac rehabilitation, 343
heart rate profile during: and sudden death, 386
intensity of
comparison of Counting Talk Test with heart rate measures and Bork's Rating of Perceived Exertion for estimating, 309
music, percussion, and metronome rhythms and, 309
Talk Test vs oxygen consumption definition of, 308
at ventilatory threshold: %HRmax and, 307
outcomes after pulmonary rehabilitation, in COPD, 186
and oxidative stress, 220
postexercise hypotension: exercise duration and, 366
and postprandial metabolism and vascular function, in men, 123
prescription for, after myocardial infarction, 361
risk-protection paradox of, 189
tolerance for
with combination tiotropium and pulmonary rehabilitation, in COPD, 386
and enrollment in cardiac rehabilitation, 343
treadmill: criterion-related validation of ventilatory response index for, 307
Exercise electrocardiography: diagnostic value of heart rate during, 127
Exercise equipment: obstacles to use of, for persons with disabilities, 301
Exercise testing
blood glucose levels during, 222
cardiopulmonary, in heart failure, 292
maximal graded
and glutathione, 215
and glutathione, in CAD, 292
North Carolina 6-minute cycle test
comparison with 6-minute walk test, 293
validity and reliability of, 292
submaximal: prediction of maximal heart rate through, 290
Exercise training
and β-blockers, 354
for cardiac patients, 296
and dyspnea self-management, 275, 286
in metabolic syndrome and diabetes, 59
whole-body: effects on body composition and functional capacity, in COPD, 124

Fairview Health System: Transtheoretical Model-based staff training and clinical intervention, 301
Family functioning: and medical adherence in cardiac rehabilitation, 300
Fatigue, skeletal muscle: aerobic-resistance training and, in men with coronary heart disease, 312
Fatty acids, polyunsaturated: and exercise capacity in COPD, 237
Fitness: in obesity, metabolic syndrome, and diabetes mellitus, 59
Frequently asked questions, 67
Functional deficits
and enrollment in cardiac rehabilitation, 343
pain or difficulty ratings as indicators of, after coronary artery bypass, 300

Gender differences
in C-reactive protein and cholesterol, 304
in clinical responsiveness to therapeutic lifestyle changes, 300
in quality of life outcomes in elderly cardiac rehabilitation patients, 295
in risk factor status and completion of phase 2 cardiac rehabilitation, 304
in risk factor status on entry into cardiac rehabilitation, 297
Geographic distribution of cardiac rehabilitation programs
problem of, 85
in United States, 80
Glucose blood levels, during exercise stress testing, 222
Glutathione
maximal graded exercise test and, 215
maximal graded exercise test and, in CAD, 292

Hand-held dynamometry: strength assessment with, 266
Health behavior change: enhancing patient motivation for, 207
Health markers: association with pedometry, 298
Health-related quality of life
adherence to cardiac rehabilitation and, 291
distractive auditory stimuli during upper extremity training and, in COPD, 50
Heart attack
symptoms of: knowledge of, in older men and women at risk for acute MI, 33
warning signs for, 40
Heart disease
comparison of diets for risk reduction for, 184
coronary
aerobic-resistance training and skeletal muscle fatigue in men with, 312
causal attribution for, in female patients, 135
women's perceptions of, 146
ischemic
cardiovascular risk markers in: exercise and stress management training and, 235
stress management for women with, 93
life-years gained from risk factor changes in England and Wales, 185
physical activity regulation in, motivational model of, 312
Heart failure
acute exercise and quality of life in, 311
behavioral management of: qualitative research on, 164

cardiopulmonary exercise testing in, 292
chronic
relaxation response randomized trial of, 149
resting end-tidal carbon dioxide in, reliability of, 177
diastolic: pedometer steps in, 293
lived experience of adherence for patients with, 158
one repetition maximum strength testing in, 306
systolic: pedometer steps in, 293
Heart rate
diagnostic value of, 127
during exercise: and sudden death, 386
horticultural therapy and, 270
maximal (HR_{max})
comparison with Counting Talk Test, 309
and exercise intensity at ventilatory threshold, 307
prediction through submaximal exercise tests, 290
after myocardial infarction or coronary artery bypass graft surgery: controlled breathing program and, 308
recovery after treadmill stress testing: depression and, 307
Heart rate reserve, 309
Home-based cardiac rehabilitation
patient preferences for, 24
as preferred and viable option, 30
Home-based pulmonary rehabilitation: in COPD, 378
Home-based walking and strength training program for obese elderly with osteoarthritis of knee, 309
Horticultural therapy: effects on mood and heart rate, 270
Hospital-based cardiac rehabilitation: patient preferences for, 24
Hospitalization for acute exacerbations of COPD: community pulmonary rehabilitation after, 186
Human Activity Profile, 115
Hypotension, postexercise: exercise duration and, 366

In memoriam: Michael S. Stulbarg, 285
Injury, spinal cord: physical activity in, 299
Insulin: capillary vs venous responses, 305
Interviewing, motivational: and secondary prevention, in cardiac rehabilitation, 200
Ischemic heart disease
cardiovascular risk markers in: exercise and stress management training effects on, 235
stress management for women with, 93

Journal of Cardiac Rehabilitation, 243

Knee osteoarthritis: home-based walking and strength training program for obese elderly with, 309

Life expectancy: life-years gained from cardiological treatments and risk factor changes in England and Wales, 185
Lifestyle modification
multicomponent: adherence to, 291
therapeutic: gender differences in clinical responsiveness to, 300
Lipids
cardiac rehabilitation and, 257, 264
outcomes one year after cardiac rehabilitation, 295
Literature Update, 56, 122, 184, 235, 386
Lived experience of adherence for heart failure patients, 158
Lung disease. *See* also Chronic obstructive pulmonary disease
Seattle Obstructive Lung Disease Questionnaire (SOLDQ), 107

Medical adherence: social support, family functioning, depression, and quality of life and, in cardiac rehabilitation, 300
Medical directors: responsibilities for outpatient cardiac rehabilitation/secondary prevention programs, 315
Men
cardiac rehabilitation enrollees: depressive disorders in, 71
with coronary heart disease: aerobic-resistance training and skeletal muscle fatigue in, 312
older
cardiac rehabilitation and cognitive performance in, 173
at risk for acute MI: knowledge of heart attack symptoms in, 33
prior exercise effects effects on postprandial metabolism and vascular function in, 123
Metabolic syndrome
cardiac rehabilitation and, in diabetes mellitus, 298
cardiac rehabilitation and exercise training in, 59
in COPD, 226, 233
fitness in, 59
Metabolism, postprandial: prior exercise effects on, in men, 123
Metronome rhythms: and exercise intensity, 309
Mood: horticultural therapy and, 270
Motivation for behavior change: enhancing, 207
Motivational interviewing: and secondary prevention, in cardiac rehabilitation, 200
Multicomponent lifestyle modification programs: adherence to, 291
Muscle function: and shuttle walking performance, in COPD, 43
Music
and exercise intensity, 309
during upper extremity training, in COPD, 50
Myocardial infarction
acute: knowledge of heart attack symptoms in older men and women at risk for, 33
controlled breathing program and heart rate variability after, 308
exercise intensity prescription after, 361
posttraumatic stress disorder after, 166
risk factors for, potentially modifiable, case-control study, 56
Myocardial ischemia, stress-induced: and coronary risk factors and coronary artery calcification, 305
Myocardial perfusion single-photon emission computed tomography, ischemic: and coronary artery calcification, 305

NewLeaf VO_2 testing: validation of, 306
Nomograms: prognostic value for exercise capacity in women, 386
North Carolina 6-minute cycle test
comparison with 6-minute walk test, 293
validity and reliability of, 292
North Carolina 6-minute walk test, 293
Nurses: and smoking, 198

Obesity
cardiac rehabilitation in, 62
fitness in, 59
home-based walking and strength training program for elderly with osteoarthritis of knee and, 309
Obstructive pulmonary disease. *See* also Chronic obstructive pulmonary disease
Seattle Obstructive Lung Disease Questionnaire (SOLDQ), 107
Occupational outcomes: risk factor approach to improve, in cardiac rehabilitation, 311
Older adults
cardiac rehabilitation patients
cognitive performance in, 173
quality of life outcomes for, 295
obese, with osteoarthritis of knee: home-based walking and strength training program for, 309
at risk for acute MI: knowledge of heart attack symptoms in, 33
women: physical activity and cognitive function in, 56
Ornish diet for weight loss and heart disease risk reduction, 184
Osteoarthritis of knee: home-based walking and strength training program for obese elderly with, 309

Outcome measures: quality of life, in pulmonary disease, 321
Outpatient programs
  8 week pulmonary rehabilitation: functional improvements with, 296
  cardiac rehabilitation: medical director responsibilities for, 315
Oxidative stress
  acute: maximal graded exercise test and glutathione marker, 215
  and exercise, 220
Oxygen consumption
  comparison with Counting Talk Test, 308
  and exercise intensity, 308
Oxygen saturation (VO_2)
  maximum: experimental test of, 307
  NewLeaf testing of, validation of, 306
  pre and post predicted VO2 measurements in phase II cardiac rehabilitation, 303
Oxygen therapy, long-term: and daily activity profile in COPD, 181
Oxygen uptake on-kinetics, short-term reliability of, 306

Pain ratings: and functional deficits after coronary artery bypass, 300
Patient preferences for home-based versus hospital-based cardiac rehabilitation, 24
Pedometry
  association with health markers, 298
  in diastolic and systolic heart failure, 293
Perceived control: and PTSD after myocardial infarction, 166
Perceived exertion: Bork's Rating of, 309
Percussion: and exercise intensity, 309
Peripheral muscle function: and shuttle walking performance, in COPD, 43
$P_{ET}CO_2$. *See* Resting end-tidal carbon dioxide
Physical activity
  and cognitive function, in older women, 56
  after hospitalization for CAD, 311
  regulation in heart disease, motivational model of, 312
  2 years after cardiac rehabilitation, 14
Physical exercise. *See* also Exercise
  anti-inflammatory effects of, 339
Physical functioning: cognitive performance and, after coronary artery bypass, 300
Pilates training: physiological responses to, 308
Polyunsaturated fatty acids: and exercise capacity in COPD, 237
Posttraumatic stress disorder: after myocardial infarction, 166
Posture, sitting: effects on pulmonary and cardiovascular function in COPD, 305
Prevention, secondary
  in cardiac rehabilitation: motivational interviewing and, 200
  medical director responsibilities for programs, 315
Psychosocial functioning: cognitive performance and, after coronary artery bypass, 300
Psychosocial Risk Factor Survey—PRFS, 301
PTSD. *See* Posttraumatic stress disorder
PUFAs. *See* Polyunsaturated fatty acids
Pulmonary disease. *See* Chronic obstructive pulmonary disease
Pulmonary fibrosis, idiopathic: quadriceps weakness and exercise capacity in, 386
Pulmonary function: sitting posture and, in COPD, 305
Pulmonary rehabilitation
  8 week outpatient program: functional improvements with, 296
  combined with tiotropium, in COPD, 386
  community: after hospitalization for acute exacerbations of COPD, 186
  exercise outcomes after, in COPD, 186
  first 20 years, 243
  home-based program, in COPD, 378

Quadriceps weakness: and exercise capacity in idiopathic pulmonary fibrosis, 386

Qualitative research on adherence, 164
Quality indicators, 249
Quality of life
  acute exercise and, in heart failure, 311
  age and, in cardiac rehabilitation, 304
  in elderly cardiac rehabilitation patients, 295
  health-related
    adherence to cardiac rehabilitation and, 291
    distractive auditory stimuli during upper extremity training and, in COPD, 50
  medical adherence and, in cardiac rehabilitation, 300
  phase II cardiac rehabilitation and, 296
  PTSD after myocardial infarction and, 166
  in pulmonary disease, 321

Racial differences in treatment and outcomes of high-risk non-ST elevation acute coronary syndromes, 236
Rehabilitation. *See* Cardiac rehabilitation; Pulmonary rehabilitation
Rehabilitation equipment: obstacles to use of, for persons with disabilities, 301
Relaxation response, in chronic heart failure, 149
Religiosity: and spouse's perception of quality of life and confidence, 299
Research, qualitative, on adherence, 164
Resistance training. *See* Aerobic-resistance training
Resting end-tidal carbon dioxide ($P_{ET}CO2$): reliability in chronic heart failure, 177
Risk factor status
  and completion of phase 2 cardiac rehabilitation, 304
  coronary: stress-induced myocardial ischemia and, 305
  on entry into cardiac rehabilitation, 297

Screening for depression
  in cardiac rehabilitation: AACVPR position statement on, 11
  in phase II cardiac rehabilitation, 299
Seattle Obstructive Lung Disease Questionnaire (SOLDQ), 107
Secondary prevention
  in cardiac rehabilitation: motivational interviewing and, 200
  medical director responsibilities for programs, 315
Self-management, dyspnea: exercise training and, 275, 286
Short Form Health Survey (SF-36): elderly cardiac rehabilitation patient assessment, 295
Shortness of Breath Questionnaire (UCSD): minimal clinically important difference for, 370
Shuttle walking performance: muscle function and, in COPD, 43
Single-photon emission computed tomography (SPECT)
  ischemic myocardial perfusion: and coronary artery calcification, 305
  screening stress imaging: identification of high-risk asymptomatic diabetic candidates for, 122
Sitting posture: and pulmonary and cardiovascular function in COPD, 305
  6-minute cycle test
    comparison with 6-minute walk test, 293
    validity and reliability of, 292
  6-minute walk test: comparison with 6-minute cycle test, 293
Skeletal muscle
  fatigue: aerobic-resistance training and, in men with coronary heart disease, 312
  oxygen saturation: incremental treadmill walking and, in COPD, 298
Smoking: nurses and, 198
Social support: and medical adherence in cardiac rehabilitation, 300
SOLDQ (Seattle Obstructive Lung Disease Questionnaire), 107
Spinal cord injury: physical activity in, 299
Spouse caregivers: spirituality and religiosity and perception of quality of life and confidence, 299
Sprituality: and spouse's perception of quality of life and confidence, 299
ST-segment depression post CABG, during phase II cardiac rehabilitation, 297

Staff training, Transtheoretical Model-based. in cardiac rehabilitation, 301
Strength testing
with hand-held dynamometry, 266
one repetition maximum, in heart failure, 306
Strength training
in COPD patients, 123
home-based, for obese elderly with osteoarthritis of knee, 309
Stress
oxidative
acute, maximal graded exercise test and glutathione marker of, 215
and exercise, 220
posttraumatic stress disorder after myocardial infarction, 166
Stress-induced myocardial ischemia: and coronary risk factors and coronary artery calcification, 305
Stress management
and cardiovascular risk markers, in ischemic heart disease, 235
for women with ischemic heart disease, 93
Stress SPECT imaging, screening: identification of high-risk asymptomatic diabetic candidates for, 122
Stress testing
blood glucose levels during, 222
treadmill: depression and heart rate recovery after, 307
Stroke: warning signs for, 40
Stulbarg, Michael S.: in memoriam, 285
Sudden death: heart rate profile during exercise and, 386
Sulfonylurea: and blood glucose levels during exercise stress testing, 222
Surgery, coronary bypass
cognitive performance and psychosocial and physical functioning after, 300
controlled breathing program and heart rate variability after, 308
off-pump versus on-pump cardiac rehabilitation outcomes for, 294
pain or difficulty ratings as indicators of functional deficits after, 300
ST-segment depression after, during phase II cardiac rehabilitation, 297

Talk Test. *See* Counting Talk Test
TEACH Study. *See* Tracking Exercise After Cardiac Hospitalization (TEACH) Study
Telehealth
cardiac rehabilitation via, 302
Cardiovascular Health Best Practice Project, 311
Telehealth Program pilot (Texas State University), 302
Texas State University Physical Therapy Department: Telehealth Program pilot, 302
Tiotropium combined with pulmonary rehabilitation, in COPD, 386
Tobacco-free nurses, 198
Tracking Exercise After Cardiac Hospitalization (TEACH) Study, 311
Training
aerobic
and β-blockers, 354
women's self-efficacy after, in cardiac rehabilitation, 310
aerobic-resistance
and skeletal muscle fatigue in men with coronary heart disease, 312
women's self-efficacy after, in cardiac rehabilitation, 310
endurance, in COPD patients, 123
exercise
and β-blockers, 354
and dyspnea self-management, 275, 286
in metabolic syndrome and diabetes, 59
whole-body: effects on body composition and functional capacity, in COPD, 124
Pilates: physiological responses to, 308
strength
in COPD patients, 123
home-based, for obese elderly with osteoarthritis of knee, 309
stress management: and cardiovascular risk markers in ischemic heart disease, 235
upper extremity: distractive auditory stimuli on, in COPD, 50
ventilatory support during, in severe chronic airway obstruction, 236
yoga: physiological effects of, 290, 308
Transtheoretical Model: staff training and clinical intervention based on, in cardiac rehabilitation, 301
Treadmill stress testing
comparison of Bruce and modified Bruce protocols, 290
depression and heart rate recovery after, 307
incremental walking: effects on skeletal muscle oxygen saturation in COPD, 298
ventilatory response index for, criterion-related validation of, 307

United States: geographic distribution of cardiac rehabilitation programs in, 80
University of California San Diego (UCSD) Shortness of Breath Questionnaire: minimal clinically important difference for, 370
Upper extremity training: distractive auditory stimuli on, in COPD, 50

Vascular function, postprandial: prior exercise effects on, in men, 123
Venous blood insulin response, 305
Ventilatory response index for treadmill exercise: criterion-related validation of, 307
Ventilatory support during training, in severe chronic airway obstruction, 236
Ventilatory threshold: %HRmax and exercise intensity at, 307

Wales: life-years gained from cardiological treatments and risk factor changes in, 185
Walking
and cognitive function, in older women, 56
home-based program for obese elderly with osteoarthritis of knee, 309
muscle function and, in COPD, 43
North Carolina 6-minute walk test, 293
treadmill, incremental: effects on skeletal muscle oxygen saturation, in COPD, 298
Weakness, quadriceps: and exercise capacity in idiopathic pulmonary fibrosis, 386
Weight loss
in cardiovascular rehabilitation and maintenance, 297
comparison of diets for, 184
Weight Watchers: for weight loss and heart disease risk reduction, 184
Whole-body exercise training: effects on body composition and functional capacity, in COPD, 124
Women
in cardiac rehabilitation: depressive disorders in, 71
causal attribution for coronary heart disease in, 135
with ischemic heart disease: stress management for, 93
older
cardiac rehabilitation and cognitive performance in, 173
physical activity and cognitive function in, 56
at risk for acute MI: knowledge of heart attack symptoms in, 33
perceptions of coronary heart disease, 146
prognostic value of nomogram for exercise capacity in, 386
self-efficacy after aerobic or aerobic-resistance training, in cardiac rehabilitation, 310

Yoga training: physiological effects of, 290, 308

Zone diet for weight loss and heart disease risk reduction, 184